

A Hopf algebra having a separable Galois extension is finite dimensional

J. Cuadra*

Universidad de Almería

Depto. Álgebra y Análisis Matemático

E-04120 Almería, Spain

email: jcdiaz@ual.es

2000 Mathematics Subject Classification: Primary 16W30.

Abstract

It is shown that a Hopf algebra over a field admitting a Galois extension separable over its subalgebra of coinvariants is of finite dimension. This answers in the affirmative a question posed by Beattie et al. in [*Proc. Amer. Math. Soc.* 128 No. 11 (2000), 3201-3203].

Introduction

The notion of Hopf-Galois extension, as known nowadays, is due to Kreimer and Takeuchi [5] and it is a mainstay of Hopf algebra theory. It emanated from the work of Chase and Sweedler about actions of Hopf algebras on rings [2]. When the Hopf algebra is the coordinate algebra of an affine group scheme that acts on an affine algebra, the above notion may be interpreted in geometric terms and it is linked with the concept of torsor or principal homogeneous space [7, page 168]. Faithfully flat Hopf-Galois extensions are currently widely accepted as a noncommutative counterpart of this geometric concept.

For $H = k[G]$, the group algebra of a group G over a field k , [6, Theorem 8.1.7] shows that an H -Galois extension is precisely a strongly graded algebra. That is, a k -algebra A admitting a decomposition $A = \bigoplus_{\sigma \in G} A_{\sigma}$ as k -vector space and satisfying $A_{\sigma}A_{\tau} = A_{\sigma\tau}$ for all $\sigma, \tau \in G$. The subalgebra

*This research was supported by project MTM2005-03227 from MCYT and FEDER.

of coinvariants of A is A_e (e the identity element of G). Năstăsescu et al. characterized in [4, Proposition 2.1] when the extension $A_e \subset A$ is separable. In particular, they found that if A is separable over A_e , then G is finite. It was investigated in [1] if an analogous result could hold for general Hopf algebras. To be more precise, suppose that H is a Hopf algebra over k having a Hopf-Galois extension separable over its subalgebra of coinvariants. Is H necessarily of finite dimensional? A positive response was given under the additional assumption that H to be co-Frobenius.

In this short note we answer this question in the affirmative. Our proof relies on a combination of the properties of the separability idempotent, the Galois maps and an old result of Sweedler. As often happens in Hopf algebra theory, the new proof seems more natural and simpler than the original proof for $H = k[G]$. Combining our result with one of Cohen and Fischman we provide a characterization of separable Hopf-Galois extensions that generalizes to Hopf algebras the above-mentioned one of Năstăsescu et al. for strongly graded rings.

We fix some notation and recall the definition of Hopf-Galois extension. We expect that the reader is familiar with the rudiments of Hopf algebra theory. Our conventions and notations are those of [6]. Throughout H stands for a Hopf algebra over a field k . Its counit is denoted as usual by ε . All vector spaces considered in the sequel are over k , map means linear map, and $\otimes$ denotes the tensor product over k .

For a right H -comodule algebra A with structure map $\rho : A \rightarrow A \otimes H$, its subalgebra of coinvariants $A^{\text{co}(H)} = \{a \in A : \rho(a) = a \otimes 1_H\}$ is denoted by B . The Galois maps are given by

$$\begin{aligned} \text{can} : A \otimes_B A &\rightarrow A \otimes H, & a \otimes_B a' &\mapsto \sum_{(a')} aa'_{(0)} \otimes a'_{(1)}, \\ \text{can}' : A \otimes_B A &\rightarrow A \otimes H, & a \otimes_B a' &\mapsto \sum_{(a)} a_{(0)}a' \otimes a_{(1)}. \end{aligned}$$

Recall from [6, Definition 8.1.1] that $B \subset A$ is said to be an H -Galois extension if can is an isomorphism. It is known that for H having bijective antipode, any of the two maps to be isomorphism may be required in the definition of Galois extension since can is bijective if and only if can' is bijective, [6, page 124]. However, we will not employ this fact. Furthermore, *we will only use that can is surjective.*

1 The main theorem

Theorem. *Let H be a Hopf algebra and let A be a right H -Galois extension separable over its subalgebra of coinvariants B . Then H is finite dimensional.*

Proof: We will prove that H has a non-zero finite dimensional left ideal. In virtue of [8, Corollary 2.7] this will imply that H is finite dimensional.

Let $e = \sum_{i=1}^n e_i \otimes_B e'_i \in A \otimes_B A$ be the separability idempotent given by hypothesis. Then $\sum_{i=1}^n e_i e'_i = 1_A$ and $ae = ea$ for all $a \in A$. Notice that $can'(e)$ is non-zero since

$$1_A = \sum_{i=1}^n e_i e'_i = \sum_{i=1}^n \sum_{(e_i)} e_{i(0)} \varepsilon(e_{i(1)}) e'_i = (id_A \otimes \varepsilon) can'(e).$$

We pick non-zero elements $a_j \in A, h_j \in H$ for $j = 1, \dots, m$ such that

$$\sum_{j=1}^m a_j \otimes h_j = can'(e) = \sum_{i=1}^n \sum_{(e_i)} e_{i(0)} e'_i \otimes e_{i(1)} \quad (1)$$

and the a_j 's are linearly independent. Take $h \in H$ arbitrary. Since can is surjective we may find $c_l, d_l \in A$ for $l = 1, \dots, r$ satisfying

$$1 \otimes h = can\left(\sum_{l=1}^r c_l \otimes_B d_l\right) = \sum_{l=1}^r \sum_{(d_l)} c_l d_{l(0)} \otimes d_{l(1)}. \quad (2)$$

For $l = 1, \dots, r$ we have $d_l e = e d_l$. Applying can' to this equality we get

$$\sum_{i=1}^n \sum_{(e_i)} \sum_{(d_l)} d_{l(0)} e_{i(0)} e'_i \otimes d_{l(1)} e_{i(1)} = \sum_{i=1}^n \sum_{(e_i)} e_{i(0)} e'_i d_l \otimes e_{i(1)}. \quad (3)$$

Then,

$$\begin{aligned} \sum_{j=1}^m a_j \otimes h h_j &\stackrel{(1)(2)}{=} \sum_{l=1}^r \sum_{(d_l)} \sum_{i=1}^n \sum_{(e_i)} c_l d_{l(0)} e_{i(0)} e'_i \otimes d_{l(1)} e_{i(1)} \\ &\stackrel{(3)}{=} \sum_{l=1}^r \sum_{i=1}^n \sum_{(e_i)} c_l e_{i(0)} e'_i d_l \otimes e_{i(1)} \\ &\stackrel{(1)}{=} \sum_{l=1}^r \sum_{j=1}^m c_l a_j d_l \otimes h_j. \end{aligned}$$

Let $\varphi_t \in A^*$ be such that $\varphi_t(a_j) = \delta_{tj}$, the Kronecker symbol, for $t = 1, \dots, m$. Evaluating $\varphi_t \otimes id_H$ on the preceding set of equalities we obtain

$$hh_t = \sum_{l=1}^r \sum_{j=1}^m \varphi_t(c_l a_j d_j) h_j.$$

This yields that the subspace spanned by the h_j 's is a finite dimensional non-zero left ideal of H , as required. *q.e.d.*

Cohen and Fischman provided in [3, Theorem 1.8] several characterizations of separable Hopf-Galois extensions for a finite dimensional Hopf algebra. These characterizations together with our result allow to characterize such extensions for an arbitrary Hopf algebra.

Corollary. *Let H be a Hopf algebra and let A be a right H -Galois extension. Then, $A^{co(H)} \subset A$ is separable if and only if H is finite dimensional and one of the equivalent conditions (2)-(6) in [3, Theorem 1.8] holds.*

This corollary may be viewed as a generalization to Hopf algebras of [4, Proposition 2.1] characterizing strongly graded rings that are separable over its component of degree one.

References

- [1] M. Beattie, S. Dăscălescu and Ş. Raianu, *A Co-Frobenius Hopf Algebra with a Separable Galois Extension is Finite*. Proc. Amer. Math. Soc. **128** No. 11 (2000), 3201-3203.
- [2] S.U. Chase and M.E. Sweedler, *Hopf Algebras and Galois Theory*. Lecture Notes in Mathematics **97**. Springer-Verlag, Berlin, 1969.
- [3] M. Cohen, D. Fischman, *Semisimple Extensions and Elements of Trace 1*. J. Algebra **149** (1992), 419-437.
- [4] C. Năstăsescu, M. Van den Bergh and F. Van Oystaeyen, *Separable Functors Applied to Graded Rings*. J. Algebra **123** (1989), 397-413.
- [5] H.F. Kreimer, M. Takeuchi, *Hopf Algebras and Galois Extensions of an Algebra*. Indiana Univ. Math. J. **30** (1981), 675-692.
- [6] S. Montgomery, *Hopf Algebras and Their Actions on Rings*. CMBS No. 82, AMS, 1993.

- [7] H.-J. Schneider, *Principal Homogeneous Spaces for Arbitrary Hopf algebras*. Israel J. Math. **72** Nos. 1-2 (1980), 167-195.
- [8] M.E. Sweedler, *Integrals for Hopf Algebras*. Ann. of Math. (2) **89** (1969), 323-335.